

University of Groningen

Waist-to-height ratio, waist circumference and BMI as indicators of percentage fat mass and cardiometabolic risk factors in children aged 3-7 years

Sijtsma, Anna; Bocca, Gianni; L'abée, Carianne; Liem, Eryn T; Sauer, Pieter J J; Corpeleijn, Eva

Published in:
Clinical Nutrition

DOI:
[10.1016/j.clnu.2013.05.010](https://doi.org/10.1016/j.clnu.2013.05.010)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2014

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Sijtsma, A., Bocca, G., L'abée, C., Liem, E. T., Sauer, P. J. J., & Corpeleijn, E. (2014). Waist-to-height ratio, waist circumference and BMI as indicators of percentage fat mass and cardiometabolic risk factors in children aged 3-7 years. *Clinical Nutrition*, 33(2), 311-315. <https://doi.org/10.1016/j.clnu.2013.05.010>

Copyright


Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Original article

Waist-to-height ratio, waist circumference and BMI as indicators of percentage fat mass and cardiometabolic risk factors in children aged 3–7 years



Anna Sijtsma^a, Gianni Bocca^b, Carianne L'Abée^b, Eryn T. Liem^b, Pieter J.J. Sauer^b,
Eva Corpeleijn^{a,*}

^a Department of Epidemiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

^b Department of Pediatrics, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

ARTICLE INFO

Article history:

Received 8 January 2013

Accepted 17 May 2013

Keywords:

Anthropometry
Body composition
Blood pressure
Lipid
Insulin
Cholesterol

SUMMARY

Objective: To assess whether waist-to-height-ratio (WHtR) is a better estimate of body fat percentage (BF %) and a better indicator of cardiometabolic risk factors than BMI or waist circumference (WC) in young children.

Methods: WHtR, WC and BMI were measured by trained staff according to standardized procedures. ²H₂O and ²H₂¹⁸O isotope dilution were used to assess BF% in 61 children (3–7 years) from the general population, and bioelectrical impedance (Horlick equation) was used to assess BF% in 75 overweight/obese children (3–5 years). Cardiometabolic risk factors, including diastolic and systolic blood pressure, HOMA2-IR, leptin, adiponectin, triglycerides, total cholesterol, HDL- and LDL-cholesterol, TNFα and IL-6 were determined in the overweight/obese children.

Results: In the children from the general population, after adjustments for age and gender, BMI had the highest explained variance for BF% compared to WC and WHtR ($R^2 = 0.32, 0.31$ and 0.23 , respectively). In the overweight/obese children, BMI and WC had a higher explained variance for BF% compared to WHtR ($R^2 = 0.68, 0.70$ and 0.50 , respectively). In the overweight/obese children, WHtR, WC and BMI were all significantly positively correlated with systolic blood pressure ($r = 0.23, 0.30, 0.36$, respectively), HOMA2-IR ($r = 0.53, 0.62, 0.63$, respectively), leptin ($r = 0.70, 0.77, 0.78$, respectively) and triglycerides ($r = 0.33, 0.36, 0.24$, respectively), but not consistently with other parameters.

Conclusion: In young children, WHtR is not superior to WC or BMI in estimating BF%, nor is WHtR better correlated with cardiometabolic risk factors than WC or BMI in overweight/obese children. These data do not support the use of WHtR in young children.

© 2013 Elsevier Ltd and European Society for Clinical Nutrition and Metabolism. All rights reserved.

1. Introduction

Various measures are used to detect obesity – defined in terms of excess body fat – and the risk of obesity-related co-morbidities. Body mass index (BMI) is the most commonly used measure, but waist circumference (WC) and waist-to-height ratio (WHtR) as measures of abdominal fat are also used. WHtR may have an advantage over BMI because BMI provides no information about body fat distribution, in particular abdominal fat. Central fat distribution is associated with greater health risks than total body

fat.^{1,2} Therefore, WHtR may be a better indicator of cardiometabolic risk factors than BMI. In adults, WHtR is found to be a better measure than BMI and WC for the prediction of obesity-related cardiometabolic risks factors.^{3,4} An advantage of WHtR over BMI and WC in adults is that a general cut-off value of 0.5 can be used for both men and women across many ethnicities.^{3,4} Moreover, the advantage of WHtR over WC is that WHtR adjusts for height. When compared to short people with the same WC, tall people have lower levels of cardiometabolic risk factors and a 30% lower prevalence of the metabolic syndrome.⁵

WHtR decreases from birth to the age of five from 0.69 to 0.48,⁶ and this decrease continues until early adolescence to 0.40–0.41.⁷ From then on it increases slightly to 0.42–0.43 towards the age of 18. Therefore, one cut-off value for all ages during childhood and adolescence is not feasible.^{6,7} In contrast to adults, it is less clear in young children whether WHtR is better than BMI or WC in

* Corresponding author. Department of Epidemiology (FA40), University Medical Center Groningen, PO Box 30.001, 9700 RB Groningen, The Netherlands. Tel.: +31 050 3610738; fax: +31 050 3614493.

E-mail address: e.corpeleijn@umcg.nl (E. Corpeleijn).

predicting obesity-related cardiometabolic risk factors. Most of the studies available evaluated the relationship between WHtR and body fat percentage (BF%) or cardiometabolic risk factors over a large age range, including both children and adolescents, with most studies using children aged 6 years or older. To prevent overweight and obesity in adolescents and adults, early detection of overweight and obesity in young children is needed. During early childhood especially (2–6 years of age), excessive BMI gain is predictive of obesity and cardiometabolic risk in adolescence⁸ and adulthood.^{9–11} Furthermore, adipose tissue produces cytokines, like IL-6 and TNF α , which may cause metabolic syndrome, this is already seen in children.¹² These cytokines may be a possible link between insulin resistance and adiposity. Increased levels of the separate components of metabolic syndrome have already been demonstrated in children 6–9 years old.¹³ In children aged younger than 6 it is unknown whether these processes are present yet.

Few studies have examined the association between WHtR and BF%^{14,15} or cardiometabolic risk factors^{15–17} in children aged younger than 6 years. Of the three studies investigating the associations between WHtR and cardiometabolic risk factors in children aged younger than 6 years, only one study analysed the associations between WHtR and more than one cardiometabolic risk factor. Moreover, these studies lack information about overweight/obese children, despite these children being the target group for treatment programmes.

The aim of this study is to assess whether WHtR is a better estimate of BF% than BMI and WC in very young non-obese and obese children (3–7 years of age), and whether it is a better indicator of obesity-related cardiometabolic risk factors.

2. Materials and methods

2.1. Subjects

Three groups of children were included in this study. The first group consisted of children ($n = 30$) from the general healthy Dutch population, 3–4 years of age, who were randomly selected from the GECKO Drenthe cohort.¹⁸ Data were collected between March 2010 and March 2011. The second group consisted of children ($n = 31$) from the general healthy Dutch population, 6–7 years of age. These children were recruited through advertisements in a local newspaper and on the hospital information site, and by word of mouth. Data were collected in October 2006 and have been partly described in previous studies.^{19,20} In these two groups, children having medical problems which could affect physical activity, and children diagnosed with a disease or using medication known to affect body composition were excluded from the study. These two groups were combined as one group of children from the general population. The third group consisted of overweight/obese children ($n = 75$), 3–5 years of age, who were part of a randomized controlled clinical trial (GECKO Outpatients Clinic) aimed at reducing overweight.²¹ The baseline data of the participants were used for this analysis and were measured during the children's first visit to the hospital. Data were collected between October 2006 and March 2008. Only children with overweight or obesity, according to the International Obesity Task Force (IOTF) definitions²² were included. Children with overweight or obesity due to known medical conditions or eating disorders according to the Dutch Eating Behaviour Questionnaire, mental retardation, severe behavioural problems or other criteria interfering with participation were excluded. Almost all of the children were Caucasian, one child from the first group had an Asian father and two children had African parents (one in the second and one in the third group). Written informed consent was obtained for all children, and the studies were approved by the Medical Ethics Committee of the University Medical Center Groningen (UMCG).

2.2. Body composition

In all children, height to the nearest 0.1 cm and weight to the nearest 0.1 kg were measured without shoes in light clothing using a stadiometer and a calibrated digital scale, respectively. BMI was calculated as weight/height². WC was performed with a standard tape measure and recorded to the nearest 0.1 cm in standing position. WC was measured at the mid-point between the lower costal margin and the level of the anterior superior iliac. BF% was measured using different methods for the three groups. In the first group (3–4 years of age from the general population), BF% was determined by the ²H₂¹⁸O isotope dilution method. Children drank a weighted amount (around 50 grams) of doubly labelled water (Buchem, Apeldoorn, the Netherlands [²H₂O: 6.02%, H₂¹⁸O: 12.05%]; Campro Scientific, Berlin, Germany [²H₂O: 6.63%, H₂¹⁸O: 11.50%]). Saliva samples were collected by the parents before administration and approximately 4, 16, 72 and 120 h after administration. Total body water (TBW) was determined from the dilution spaces of both isotopes. The component TBW in fat free mass (FFM) was set at 0.775 for the 3-year-old boys, 0.770 for the 4-year-old boys, 0.779 for the 3-year-old girls, and 0.777 for the 4-year-old girls.²³ BF% was calculated as $([\text{weight} - \text{FFM}]/\text{weight}) \times 100$. More detailed information is described elsewhere.²⁴ In the second group (6–7 years of age from the general population), BF% was calculated from TBW, determined by an orally administered dose of 99.8% ²H₂O of 0.15 g per kilogram body weight. Detailed information about the method has been described before.²⁰ In the third group (3–5 years of age overweight/obese children), BF% was estimated using a 50 kHz fixed frequency hand-to-foot bio-impedance analyzer (BIA-101, Akern S.r.l./RJL Systems, Florence, Italy). The measurements were performed three times and the average calculated. The resistance (R_z) value from BIA together with height, weight, age and sex were used to calculate the FFM using the Horlick equation.²⁵ BF% was calculated as $([\text{weight} - \text{FFM}]/\text{weight}) \times 100$.

2.3. Overweight/obesity-related cardiometabolic risk factors

Cardiometabolic risk factors were only measured in the overweight/obese group. Systolic and diastolic blood pressures were measured in supine position at the right upper arm with a Dinamap Critikon 1846SX digital sphygmomanometer (Critikon Inc., Tampa, Florida, USA) and an appropriate cuff size. The child was instructed not to speak or move during the measurements. The mean of two measurements was calculated. Blood was drawn after an overnight fast. An enzymatic colorimetric method (Roche Modular, Basel, Switzerland) was used to determine total cholesterol, high-density lipoprotein (HDL)-cholesterol, low-density lipoprotein (LDL)-cholesterol and triglycerides. The updated homeostasis model assessment of insulin resistance (HOMA2-IR) was used to calculate insulin resistance.²⁶ HOMA2-IR was calculated from fasting plasma glucose and fasting plasma insulin, determined by enzymatic method (hexokinase-mediated reaction, Roche Modular, Basel, Switzerland) and radioimmunoassay (Diagnostic Systems Laboratories, Inc., Webster, TX, USA), respectively. Serum levels of leptin, interleukin-6 (IL-6) and tumour necrosis factor- α (TNF α) were measured by a combination enzyme-linked immunosorbent assay (ELISA; Milliplex Map Human Adipokine Panel B, Millipore, St. Charles, MN, USA). Serum levels of adiponectin were quantified by ELISA (Millipore, St. Charles, MN, USA).

2.4. Statistics

Mean and standard deviations were calculated for all of the characteristics measured. Univariate linear regression analyses were performed with BF% as the dependent variable and WHtR, WC

Table 1
Anthropometric characteristics in the three groups of children.

	3–4 years of age General population (<i>n</i> = 30, 40% boys)		6–7 years of age General population (<i>n</i> = 31, 56% boys)		3–5 years of age Overweight/obese population (<i>n</i> = 75, 28% boys)	
	Mean	SD	Mean	SD	Mean	SD
Age	3.5	0.3	6.7	0.5	4.7	0.8
Weight (kg)	16.3	1.9	25.1	3.9	28.2	6.5
Height (cm)	101.0	5.4	124.8	6.4	114.9	7.8
Waist (cm)	51.7	2.7	57.0	3.9	64.9	7.5
BMI	15.9	1.2	16.0	1.6	21.1	2.8
WHTR	0.51	0.03	0.46	0.03	0.56	0.05
BF% ^a	22.0 ^b	6.4	17.5 ^b	5.3	28.3 ^c	6.8

BF%, body fat percentage; BMI, body mass index; WC, waist circumference; WHTR, waist-to-height ratio.

^a Data on BF% were missing in four children aged 3–4 years, in three children aged 6–7 years, and in nine overweight/obese children aged 3–5 years.

^b BF% as determined by BIA with Horlick equation.

^c BF% as determined by isotope dilution method.

or BMI as the independent factors. In the multivariate linear regression analyses gender and age were added as independent factors. Effect modification of age or gender with WHTR, WC and BMI were checked. Pearson correlations were used to analyse the relationship between the BF% or cardiometabolic risk factors and WHTR, WC or BMI. Statistical analyses were performed using PASW 18.0.3 for Windows (SPSS, Chicago Illinois, USA). The significance level was set to $p < 0.05$ (2-tailed).

3. Results

Table 1 shows the characteristics of the three groups of children. The children aged 3–4 years ($n = 30$) and 6–7 years ($n = 31$) were representatives of the general population, which was reflected in the number of overweight children, three and six children were overweight, based on the IOTF definitions,²² respectively. The third group consisted of overweight/obese children aged 3–5 years ($n = 75$). Overweight was present in 38.7% ($n = 29$) of the children

and 61.3% ($n = 46$) were obese. In addition to the anthropometric measures, the following cardiometabolic measures were also determined in the overweight/obese children: systolic and diastolic blood pressure (113 ± 13 and 64 ± 7 mmHg, respectively), HOMA2-IR (1.0 ± 0.5), leptin (9.4 ± 9.4 ng/mL), adiponectin (18.3 ± 4.7 ng/mL), triglycerides (69.0 ± 33.1 mg/dL), total cholesterol (149.0 ± 23.4 mg/dL), HDL-cholesterol (49.5 ± 10.4 mg/dL), LDL-cholesterol (95.8 ± 20.9 mg/dL), TNF α (3.6 ± 3.2 pg/mL) and IL-6 (1.5 ± 1.2 pg/mL).

Table 2 shows the associations of BF% with WHTR, WC and BMI. In the crude models, WHTR ($r = 0.48$, $p < 0.001$) and BMI ($r = 0.44$, $p < 0.001$) were associated with BF% in the children from the general population when age groups were combined. In the overweight/obese children, WHTR was correlated with BF% ($r = 0.65$, $p < 0.001$), but WC and BMI were even better correlated with BF% ($r = 0.81$, $p < 0.001$ and $r = 0.81$, $p < 0.001$, respectively). When adjusted for gender and age, WHTR, WC and BMI were significantly associated with BF% in all groups. In the adjusted model for the children from the general population with age groups combined, BMI had the highest explained variance ($R^2 = 0.32$) for BF%. This was also true for the children of 6–7 years of age, but WHTR performed best in the children of 3–4 years of age. In the adjusted models for the overweight/obese children, WHTR had the lowest explained variance ($R^2 = 0.50$), compared to WC ($R^2 = 0.70$) and BMI ($R^2 = 0.68$). We checked if effect modifications of age or gender with WHTR, WC or BMI were present in the models on top of the adjustments for age and gender. The only significant interaction term was for gender with WHTR in the children from the general population with age groups combined. Adding this term to the model for predicting BF%, the explained variance increased from $R^2 = 0.23$ to $R^2 = 0.33$. This resulted in a beta coefficient of 74.9 for boys and 74.0 for girls. The additional adjustment for interaction with gender makes the explained variance of WHTR, WC or BMI for BF% comparable.

Table 3 shows the associations of WHTR, WC and BMI with cardiometabolic risk factors for the overweight/obese children. WHTR, WC and BMI were positively correlated with systolic blood pressure, HOMA2-IR, leptin and triglycerides. For these significant

Table 2
Associations of WHTR, WC and BMI with body fat percentage.

	Body fat percentage											
	3–4 years of age General population ^a			6–7 years of age General population ^a			3–4 and 6–7 years of age General population ^a			3–5 years of age Overweight/obese population ^b		
	B [95% CI]	R ² model	Pearson r	B [95% CI]	R ² model	Pearson r	B [95% CI]	R ² model	Pearson r	B [95% CI]	R ² model	Pearson r
WHTR												
Crude	67.5 [−9.1;144.0]	0.12	0.35	56.2 [−9.5;121.9]	0.11	0.33	72.6 [35.7; 109.6]	0.23***	0.48***	92.9 [66.1;119.8]	0.43***	0.65***
+Gender	72.5 [−0.4;145.4]	0.24*		84.1 [20.2;147.9]	0.29*		73.1 [34.7; 111.4]	0.23***		94.5 [66.7;122.4]	0.43***	
+Gender, age	94.8 [25.7;164.0]	0.40**		85.8 [20.3;151.3]	0.30*		71.3 [20.5; 122.0]	0.23***		97.4 [71.2;123.7]	0.50***	
WC												
Crude	1.09 [0.24;1.93]	0.23*	0.48*	0.75 [0.30;1.21]	0.31**	0.56**	0.17 [−0.22; 0.56]	0.02	0.12	0.77 [0.63;0.90]	0.66***	0.81***
+Gender	1.09 [0.29;1.89]	0.34**		0.89 [0.48;1.30]	0.49***		0.17 [−0.22; 0.57]	0.03		0.80 [0.66;0.94]	0.67***	
+Gender, age	0.99 [0.14;1.84]	0.35*		0.93 [0.51;1.36]	0.51***		0.87 [0.42; 1.33]	0.31***		0.87 [0.71;1.02]	0.70***	
BMI												
Crude	1.64 [−0.42;3.70]	0.10	0.32	2.18 [1.24;3.12]	0.46***	0.68***	1.87 [0.79; 2.94]	0.19***	0.44***	2.14 [1.75;2.53]	0.65***	0.81***
+Gender	1.72 [−0.25;3.69]	0.22		2.34 [1.50;3.17]	0.61***		1.85 [0.77; 2.93]	0.20***		2.27 [1.87;2.67]	0.67***	
+Gender, age	2.14 [0.24;4.03]	0.35*		2.34 [1.49;3.20]	0.61***		1.96 [0.95; 2.97]	0.32***		2.23 [1.82;2.63]	0.68***	

BMI, body mass index; WC, waist circumference; WHTR, waist-to-height ratio.

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

^a Body fat percentage measured by isotope dilution method

^b Body fat percentage measured by BIA with Horlick equation.

Table 3

Pearson correlation coefficients regarding the associations between cardiometabolic risk factors and WHtR, WC and BMI in overweight/obese children aged 3–5 years.

	WHtR (r)	WC (r)	BMI (r)
SBP	0.233*	0.303**	0.361**
DBP	0.076	0.127	0.168
HOMA2-IR	0.534***	0.618***	0.627***
Leptin ^a	0.701***	0.769***	0.779***
Adiponectin	−0.211	−0.311**	−0.152
TG	0.330**	0.364**	0.244*
TC	0.035	0.085	0.056
HDL-C	−0.231*	−0.040	−0.012
LDL-C	0.064	0.033	0.038
TNF α	0.094	0.107	0.158
IL-6	0.030	−0.057	−0.018

BMI, body mass index; DBP, diastolic blood pressure; HDL-C, high-density lipoprotein cholesterol; HOMA2-IR, updated homeostasis model assessment of insulin resistance; IL-6, interleukin-6; LDL-C, low-density lipoprotein cholesterol; SBP, systolic blood pressure; TC, total cholesterol; TG, triglycerides; TNF α , tumour necrosis factor- α ; WC, waist circumference; WHtR, waist-to-height ratio. Less than 4% of data was missing, with the exception of HOMA2-IR, which was missing in 10 children (13%).

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

^a Log transformed.

associations, WHtR was not superior to WC and BMI. For adiponectin we found only a significantly inverse correlation with WC and for HDL-cholesterol only an inverse correlation with WHtR.

4. Discussion

This study revealed that WHtR is associated with BF% in children. However, even after adjustments for gender and age, WHtR is not better associated with BMI and WC. Neither is WHtR a better indicator of most cardiometabolic risk factors than WC and BMI.

4.1. WHtR, WC and BMI as indicators of body fat percentage

Previous studies that investigated the association between WHtR and BF% in both normal and overweight children found that WHtR was associated with total or abdominal BF%.^{15,27–31} Studies in older children (8–16 years of age) showed a better correlation of WHtR with BF% than BMI.^{27,28} Studies that only included children younger than 10 years have found that WHtR was not better correlated with BF% than BMI or WC.^{15,29,30} Therefore, this association may be dependent on age. The correlation between WHtR and BF% has been evaluated in only two studies of children of less than 6 years of age.^{14,15} In these studies the prevalence of overweight/obesity was around 30%^{14,15} and the associations were adjusted for age and shown separately for boys and girls. In both studies a significant correlation was found. In our study we also found this significant correlation, both in the overweight obese group as well as in the group from the general population, although the associations seemed to be stronger in the older children (6–7 years compared to 3 years of age). The finding that the association between WC and BF% was initially lost after combining the two age groups in one general population group stresses the importance to look at age-range specific subgroups. But, because WHtR is not better related to BF% than BMI and WC in both the unadjusted and adjusted models, we conclude that WHtR has no advantage over BMI in predicting BF%. It is common to define overweight in children based on the BMI adjusted for gender and age.²² WHtR, either with or without these adjustments, is not a better screening method than adjusted BMI and WC for predicting BF%.

We observed stronger associations between BF% and WHtR, WC and BMI in the overweight/obese group compared to the general population group. An explanation might be that BMI is a better

measure of excess body fat in obese children compared to normal weight or overweight children, because BMI is not only dependent on fat mass but also on muscle mass,³² and the higher the fat mass, the smaller the influence of variation in muscle mass. Almost 70% of the children in the overweight/obese group were obese, which may explain why BMI has a higher association with BF% in the overweight/obese group compared to the general population groups. BF% was measured using the isotope dilution method in the general population group, while in the group with overweight/obese children it was assessed by BIA and estimated using the Horlick equation. As expected, BF% was higher in the overweight/obese group compared to the children from the general population group. Even if the BF% may have been underestimated by the BIA method,²⁰ this will not have influenced the correlations with WHtR, WC and BMI, so the difference in methods is not a likely explanation for the higher correlation between the anthropometric measurements and BF%. Moreover, the overweight/obese group consisted of more children than the group from the general population. A group with more children has more power to find a significant effect, but the correlation coefficients will not automatically be higher. Finally, the standard deviations and ranges of WHtR, WC, BMI and BF% were wider in the overweight/obese group, which means a higher chance of finding an association.

4.2. WHtR, WC and BMI as indicators of cardiometabolic risk factors

We found only three studies that investigated the relationship between WHtR and one or more cardiometabolic risk factors in children younger than 6 years.^{15–17} Corvalan et al. assessed the relationship between the anthropometric measures (e.g. WHtR, WC and BMI) and different cardiometabolic markers in 3 to 4-year-old children.¹⁵ Whitrow et al. assessed whether WHtR is a better indicator than BMI of SBP in children, 3.5 years of age.¹⁷ Campagnolo et al. assessed the accuracy of WHtR, WC and BMI in identifying children (3–4 years of age) with multiple risk factors for cardiovascular disease. No correlations between the anthropometric measures and the separate cardiovascular risk factors were found.¹⁶

We found that apart from HDL-cholesterol none of the cardiovascular risk factors were better correlated with WHtR than with BMI or WC (i.e. SBP, HOMA2-IR, leptin and triglycerides). This is in accordance with most previous studies in children aged 4–17 years, based on a review by Browning et al.³ For HDL-cholesterol we found a significant inverse association with WHtR, while no association was found with BMI or WC. This inverse association was also found in a previous review,³ while no association was found in young children.¹⁵ The lack of association between total cholesterol and IL-6 with WHtR has also been found in previous studies.^{3,15} The positive association with triglycerides is consistent with studies in older children, but was not found in young children.¹⁵ The absence of an association of WHtR with DBP, adiponectin, LDL-cholesterol and TNF α in young children has not been published before.

A major strength of our study is that the sample was composed of young children of 3–7 years of age. In addition, relationships were studied in overweight/obese children. Studies analysing these relationships in young children are limited and we found no studies that analysed these in overweight/obese children at this young age, despite these children being the target population for intervention programmes. The low number of children in the general population groups is a limitation of our study. Although for diagnosis purposes, the cross-sectional design of the study is adequate, the drawback is that it gives no information about the ability of the proxy measures to predict cardiometabolic risk factors later in life. For diagnosis of excess adiposity, cross-sectional data are sufficient, but it would be

interesting and very relevant to relate the adiposity measures to future health outcomes.

In conclusion, in young children, either from the general population or overweight/obese, WHtR was not superior to WC or BMI in estimating BF%, nor was WHtR better correlated with cardiometabolic risk factors than WC or BMI in overweight/obese children. These data do not support the use of WHtR in young children.

Conflict of interest

The authors declare there are no competing financial interests in relation to the work described.

References

- Vague P. The degree of masculine differentiation of obesities: a factor determining predisposition to diabetes, atherosclerosis, gout, and uric calculous disease. *Am J Clin Nutr* 1956;**4**:20–34.
- Despres JP. Is visceral obesity the cause of the metabolic syndrome? *Ann Med* 2006;**38**:52–63.
- Browning LM, Hsieh SD, Ashwell M. A systematic review of waist-to-height ratio as a screening tool for the prediction of cardiovascular disease and diabetes: 0.5 could be a suitable global boundary value. *Nutr Res Rev* 2010;**23**:247–69.
- Ashwell M, Hsieh S. Six reasons why the waist-to-height ratio is a rapid and effective global indicator for health risks of obesity and how its use could simplify the international public health message on obesity. *Int J Food Sci Nutr* 2005;**56**:303–7.
- Schneider HJ, Klotsche J, Silber S, Stalla GK, Wittchen HU. Measuring abdominal obesity: effects of height on distribution of cardiometabolic risk factors risk using waist circumference and waist-to-height ratio. *Diabetes Care* 2011;**34**:e7.
- Roswall J, Bergman S, Almqvist-Tangen G, Alm B, Niklasson A, Nierop AFM, et al. Population-based waist circumference and waist-to-height ratio reference values in preschool children. *Acta Paediatr* 2009;**98**:1632–6.
- Brannsether B, Roelants M, Bjerknes R, Juliusson PB. Waist circumference and waist-to-height ratio in Norwegian children 4–18 years of age: reference values and cut-off levels. *Acta Paediatr* 2011;**100**:1576–82.
- Liem ET. Periods of infancy and childhood growth affecting adiposity at age 16 years. In: *Development of overweight in adolescence: genes, growth & mood*. Rijksuniversiteit Groningen; 2010. p. 66–83.
- de Kroon ML, Renders CM, van Wouwe JP, Hirasings RA, van Buuren S. Identifying young children without overweight at high risk for adult overweight: the Terneuzen Birth Cohort. *Int J Pediatr Obes* 2011;**6**:e187–95.
- de Kroon ML, Renders CM, van Wouwe JP, van Buuren S, Hirasings RA. The Terneuzen birth cohort: BMI change between 2 and 6 years is most predictive of adult cardiometabolic risk. *PLoS One* 2010;**5**:e13966.
- De Kroon ML, Renders CM, Van Wouwe JP, Van Buuren S, Hirasings RA. The Terneuzen birth cohort: BMI changes between 2 and 6 years correlate strongest with adult overweight. *PLoS One* 2010;**5**:e9155.
- Roth CL, Reinehr T. Roles of gastrointestinal and adipose tissue peptides in childhood obesity and changes after weight loss due to lifestyle intervention. *Arch Pediatr Adolesc Med* 2010;**164**:131–8.
- Martos R, Valle M, Morales RM, Canete R, Gascon F, Urbano MM. Changes in body mass index are associated with changes in inflammatory and endothelial dysfunction biomarkers in obese prepubertal children after 9 months of body mass index SD score loss. *Metabolism* 2009;**58**:1153–60.
- Taylor RW, Williams SM, Grant AM, Ferguson E, Taylor BJ, Goulding A. Waist circumference as a measure of trunk fat mass in children aged 3 to 5 years. *Int J Pediatr Obes* 2008;**3**:226–33.
- Corvalan C, Uauy R, Kain J, Martorell R. Obesity indicators and cardiometabolic status in 4-y-old children. *Am J Clin Nutr* 2010;**91**:166–74.
- Bo Campagnolo PD, Hoffman DJ, Vitolo MR. Waist-to-height ratio as a screening tool for children with risk factors for cardiovascular disease. *Ann Hum Biol* 2011;**38**:265–70.
- Whitrow MJ, Moore VM, Davies MJ. Waist-to-height ratio is not a predictor of systolic blood pressure in 3-year-old children. *J Pediatr* 2011;**159**:501–3.
- L'Abée C, Sauer PJJ, Damen M, — Rake J, Cats H, Stolk RP. Cohort profile: the GECKO Drenthe study, overweight programming during early childhood. *Int J Epidemiol* 2008;**37**:486–9.
- Liem ET, Rolfe ED, L'Abée C, Sauer PJJ, Ong KK, Stolk RP. Measuring abdominal adiposity in 6 to 7-year-old children. *Eur J Clin Nutr* 2009;**63**:835–41.
- L'Abée C, Visser GH, Liem ET, Kok DEG, Sauer PJJ, Stolk RP. Comparison of methods to assess body fat in non-obese six to seven-year-old children. *Clin Nutr* 2010;**29**:317–22.
- Bocca G, Corpeleijn E, Stolk RP, Sauer PJJ. Results of a multidisciplinary treatment program in 3- to 5-year-old children with overweight or obesity. A randomized, controlled clinical trial. *Arch Pediatr Adolesc Med* 2012;**166**:1109–15.
- Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *Br Med J* 2000;**320**:1240–3.
- Fomon SJ, Haschke F, Ziegler EE, Nelson SE. Body composition of reference children from birth to age 10 years. *Am J Clin Nutr* 1982;**35**:1169–75.
- Rieken R, Schierbeek H, van Goudoever JB, Tibboel D, Evenhuis HM, Penning C. Measuring energy expenditure and body composition in children with neurological impairment and intellectual disability using the doubly labeled water method: comparing urine and saliva sampling. *e-SPEN* 2010;**5**:e225–9.
- Horlick M, Arpad SM, Bethel J, Wang J, Moye Jr J, Cuff P, et al. Bioelectrical impedance analysis models for prediction of total body water and fat-free mass in healthy and HIV-infected children and adolescents. *Am J Clin Nutr* 2002;**76**:991–9.
- Wallace TM, Levy JC, Matthews DR. Use and abuse of HOMA modeling. *Diabetes Care* 2004;**27**:1487–95.
- Nambiar S, Hughes I, Davies PS. Developing waist-to-height ratio cut-offs to define overweight and obesity in children and adolescents. *Public Health Nutr* 2010;**13**:1566–74.
- Fujita Y, Kouda K, Nakamura H, Iki M. Cut-off values of body mass index, waist circumference, and waist-to-height ratio to identify excess abdominal fat: population-based screening of Japanese school children. *J Epidemiol* 2011;**21**:191–6.
- Maximova K, Chiolerio A, O'Loughlin J, Tremblay A, Lambert M, Paradis G. Ability of different adiposity indicators to identify children with elevated blood pressure. *J Hypertens* 2011;**29**:2075–83.
- Hubert H, Guinhouya CB, Allard L, Durocher A. Comparison of the diagnostic quality of body mass index, waist circumference and waist-to-height ratio in screening skinfold-determined obesity among children. *J Sci Med Sport* 2009;**12**:449–51.
- Taylor RW, Williams SM, Grant AM, Taylor BJ, Goulding A. Predictive ability of waist-to-height in relation to adiposity in children is not improved with age and sex-specific values. *Obesity (Silver Spring)* 2011;**19**:1062–8.
- Freedman DS, Sherry B. The validity of BMI as an indicator of body fatness and risk among children. *Pediatrics* 2009;**124**:S23–34.